

Laparoscopic Liver Resection: A South American Experience with 2887 Cases

**J. Pekolj, R. Clariá Sánchez, J. Salceda,
R. J. Maurette, P. B. Schelotto, L. Pierini,
E. Cánepa, M. Moro, G. Stork, N. Resio,
et al.**

World Journal of Surgery

Official Journal of the International
Society of Surgery/Société
Internationale de Chirurgie

ISSN 0364-2313

World J Surg

DOI 10.1007/s00268-020-05646-4



Your article is protected by copyright and all rights are held exclusively by Société Internationale de Chirurgie. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



Laparoscopic Liver Resection: A South American Experience with 2887 Cases

J. Pekolj¹ · R. Clariá Sánchez¹ · J. Salceda² · R. J. Maurette³ · P. B. Schelotto⁴ · L. Pierini⁵ · E. Cánepa⁶ · M. Moro⁷ · G. Stork⁷ · N. Resio⁸ · J. Neffa⁹ · L. Mc Cormack¹⁰ · E. Quiñonez¹¹ · G. Raffin¹² · L. Obeide¹³ · D. Fernández¹⁴ · G. Pfaffen¹⁵ · C. Salas¹⁶ · M. Linzey¹⁷ · G. Schmidt¹⁸ · S. Ruiz¹⁹ · F. Alvarez²⁰ · J. Buffaliza²¹ · R. Maroni²² · O. Campi²³ · C. Bertona²⁴ · M. de Santibañes¹ · O. Mazza¹ · M. Belotto de Oliveira²⁵ · A. L. Diniz²⁶ · M. Enne de Oliveira²⁷ · M. A. Machado²⁵ · A. N. Kalil²⁸ · R. D. Pinto²⁹ · A. P. Rezende³⁰ · E. J. B. Ramos³¹ · A. Talvane T. Oliveira³² · O. J. M. Torres³³ · N. Jarufe Cassis³⁴ · E. Buckel³⁵ · R. Quevedo Torres³⁶ · J. Chapochnick³⁷ · M. Sanhueza Garcia³⁸ · C. Muñoz³⁹ · G. Castro⁴⁰ · H. Losada⁴¹ · F. Vergara Suárez⁴² · O. Guevara⁴³ · D. Dávila⁴⁴ · O. Palacios⁴⁴ · A. Jimenez⁴⁵ · L. Poggi⁴⁶ · V. Torres⁴⁷ · G. M. Fonseca⁴⁸ · J. A. P. Kruger⁴⁸ · F. F. Coelho⁴⁸ · L. Russo⁴⁹ · P. Herman⁴⁸

© Sociét  Internationale de Chirurgie 2020

Abstract

Background Laparoscopic liver resections (LLR) have been increasingly performed in recent years. Most of the available evidence, however, comes from specialized centers in Asia, Europe and USA. Data from South America are limited and based on single-center experiences. To date, no multicenter studies evaluated the results of LLR in South America. The aim of this study was to evaluate the experience and results with LLR in South American centers.

Methods From February to November 2019, a survey about LLR was conducted in 61 hepatobiliary centers in South America, composed by 20 questions concerning demographic characteristics, surgical data, and perioperative results.

Results Fifty-one (83.6%) centers from seven different countries answered the survey. A total of 2887 LLR were performed, as follows: Argentina (928), Brazil (1326), Chile (322), Colombia (210), Paraguay (9), Peru (75), and Uruguay (8). The first program began in 1997; however, the majority (60.7%) started after 2010. The percentage of LLR over open resections was 28.4% (4.4–84%). Of the total, 76.5% were minor hepatectomies and 23.5% major, including 266 right hepatectomies and 343 left hepatectomies. The conversion rate was 9.7%, overall morbidity 13%, and mortality 0.7%.

Conclusions This is the largest study assessing the dissemination and results of LLR in South America. It showed an increasing number of centers performing LLR with the promising perioperative results, aligned with other worldwide excellence centers.

✉ P. Herman
pherman@uol.com.br

¹ HPB Surgery Section, General Surgery Service, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina

² Hospital Ramón Santamarina, Tandil, Argentina

³ Hospital Británico, Buenos Aires, Argentina

⁴ Fundación Favaloro, Buenos Aires, Argentina

⁵ Clínica Nefrología, Clínica Uruguay, Hospital Iturraspe, Santa Fe, Argentina

⁶ Hospital Privado de Comunidad, Mar del Plata, Argentina

⁷ Hospital Italiano - Regional Sur, Bahía Blanca, Argentina

⁸ Unidad HPB Sur, General Roca, Argentina

⁹ Hospital Italiano de Mendoza, Mendoza, Argentina

¹⁰ Hospital Alemán, Buenos Aires, Argentina

¹¹ Hospital El Cruce, Buenos Aires, Argentina

¹² Hospital Argerich, Buenos Aires, Argentina

¹³ Hospital Universitario Privado, Córdoba, Argentina

Introduction

The development of minimally invasive liver surgery (MILS) was slow due several barriers to overcome. The first limit was technical; hence, the translation of conventional steps to the laparoscopic approach was needed. For instance, liver mobilization, palpation, vascular dissection and control, and parenchymal transection were steps to be adapted to the laparoscopic approach. Thus, surgeons with experience in liver surgery had to be trained in advanced laparoscopy to achieve development. Moreover, bleeding control during liver transection, gas embolism secondary to pneumoperitoneum, and oncological outcomes such as surgical margins, port site seeding, and long-term survival were obstacles of great concern in the early experience with MILS [1–4].

In the last two decades, there has been a huge evolution on MILS, and many reports emerged confirming the safety and feasibility of MILS [5–7] including complex procedures such as major hepatectomies and even graft harvesting for living donation [8–10]. Regarding oncological safety, several reports have shown similar survival rates comparing open and laparoscopic resection of colorectal liver metastasis and hepatocellular carcinoma [10–17]. In addition, port seeding has proved to be extremely rare [18–21].

Evidence supporting MILS comes from case or cohort series, retrospective comparative studies, and meta-analyses. Ciria et al. in a systematic review including 9527 laparoscopic liver resections (LLR) have shown growing safety when performed in selected patients by experienced surgeons suggesting that LLR may offer improved short-term outcomes when compared to open liver resections (OLR) [22]. Only recently, a randomized controlled trial (RCT) was published comparing the results of LLR with the open approach for colorectal liver metastasis showing less postoperative complications in the laparoscopic group [23]. Despite being the best scientific evidence for the results evaluation of MILS, RCTs are difficult to conduct in clinical practice. In this context, international registries and well-designed observational studies are appropriate ways to produce evidence supporting MILS.

In 2014, the 2nd International Consensus Conference on LLR in Japan demonstrated the progress, acceptance, and dissemination of the method worldwide [24]. However, the majority of publications on LLR come from developed countries from Asia, Europe, or USA; few studies from centers in other regions of the world are available. In South America, there are limited data from single-center studies [25–29]; however, to date, no multicenter efforts to assess the dissemination and results of LLR in South America was published.

¹⁴ Clínica Pueyrredón, Mar del Plata, Argentina

¹⁵ Sanatorio Güemes, Buenos Aires, Argentina

¹⁶ Sanatorio 9 de Julio, Santiago del Estero, Argentina, Hospital Centro de Salud, San Miguel de Tucumán, Argentina

¹⁷ Hospital Angel C. Padilla, San Miguel de Tucumán, Argentina

¹⁸ Hospital Escuela Gral. Corrientes, Argentina

¹⁹ Clínica Colón, Mar del Plata, Argentina

²⁰ Clínica Reina Fabiola, Hospital Italiano, Córdoba, Argentina

²¹ Hospital Lagommaggiore, Mendoza, Argentina

²² Hospital Papa Francisco, Salta, Argentina

²³ Clínica Regional General Pico, Santa Rosa, Argentina

²⁴ Hospital Español, Mendoza, Argentina

²⁵ Hospital Sírío Libanês, São Paulo, Brazil

²⁶ A.C. Camargo Cancer Center, São Paulo, Brazil

²⁷ Ipanema Federal Hospital, Rio de Janeiro, Brazil

²⁸ Santa Casa de Porto Alegre, Universidade Federal de Ciências da Saúde, Porto Alegre, Brazil

²⁹ Hospital Santa Catarina de Blumenau, Blumenau, Brazil

³⁰ Hospital Mater Dei, Belo Horizonte, Brazil

³¹ Hospital Nossa Senhora das Graças, Curitiba, Brazil

³² Hospital Americas Medical City Hospital Samaritano, Rio de Janeiro, Brazil

³³ Hospital Universitario HUUFMA, Hospital São Domingos, UDI Hospital, Fortaleza, Brazil

³⁴ Universidad Católica de Chile, Santiago, Chile

³⁵ Clínica Las Condes, Santiago, Chile

³⁶ Hospital de Concepción, Concepción, Chile

³⁷ Clínica Santa María, Santiago, Chile

³⁸ Hospital Sotero del Río, Santiago, Chile

³⁹ Hospital de Talca, Talca, Chile

⁴⁰ Hospital de Arica, Arica, Chile

⁴¹ Hospital de Temuco, Temuco, Chile

⁴² Clínica Vida - Fundación Colombiana de Cancerología, Medellín, Colombia

⁴³ Instituto Nacional de Cancerología, Bogotá, Colombia

⁴⁴ Clínica CES, Medellín, Colombia

⁴⁵ Hospital Clínicas, Asunción, Paraguay

⁴⁶ Clínica Anglo Americana, Lima, Peru

⁴⁷ Hospital Guillermo Almenara ESSALUD, Lima, Peru

⁴⁸ Hospital das Clínicas - University of São Paulo School of Medicine, São Paulo, Brazil

⁴⁹ Hospital Maciel, Casmu, Montevideo, Uruguay

The aim of this study was to evaluate the experience and results with LLR in South American centers.

Methods

A survey including 61 hepatobiliary groups in South America, including seven countries (Argentina, Brazil, Chile, Colombia, Paraguay, Peru, and Uruguay), was performed from February to May 2019. Contacts were done by e-mail or phone calls, and the survey was sent by e-mail, with a cover letter calling for participation.

Twenty questions as follows were presented: year of the beginning of the program; total number of liver resections; number of cases OLR and LLR; percentage of laparoscopic/open cases; conversion rate; percentage of benign and malignant cases; number of colorectal metastasis, hepatocellular carcinoma, other tumors cases; number of minor LLR; number of major LLR (resection of ≥ 3 segments); number of ALPPS (associating liver partition and portal vein ligation for staged hepatectomy) cases; number of right and left LLR; number of extended LLR; number of left lateral sectionectomies; transfusion rate, morbidity, and mortality.

All studied variables, except age, were categorical and presented as frequency (percentage).

Results

There were 51 (83.6%) answers from different centers, and seven different countries as follows: Argentina (23), Brazil (11), Chile (8), Colombia (3), Paraguay (2), Peru (2), and Uruguay (2). The results are summarized in Table 1.

The first program began in 1997 and the majority (31 programs = 60.7%) after 2010. All of the 51 groups that answered the survey perform MILS. A total of 2887 LLR were performed, as follows: Argentina (928), Brazil (1326), Chile (322), Colombia (210), Paraguay (9), Peru (75), and Uruguay (8).

From the 51 responding centers, nine were considered as high-volume centers (more than 50 total liver resections per year and more than 100 LLR performed) and 42 low-volume centers. Eleven centers (21.5%) were academical university centers, and 40 (78.5%) were peripheral/regional centers.

The median age of the patients was 44 years (28–70 years). Regarding the total number of cases per group, 31 groups (60.7%) did less than 50 cases, 11 (21.5%) did between 50 and 100 cases, six (11.7%) between 100 and 150 cases, and three groups (5.8%) did more than 200 cases (Fig. 1).

Table 1 Results of 2887 laparoscopic liver resections in South America

Variable	N (%) / % (range)
# centers	51
# cases	2887
% applicability	28.4% (4.4–84)
% benign disease	43% (0–70)
% malignant tumors	57% (0–100)
# minor resections	2209 (76.5%)
# major resections	678 (23.5%)
# right hepatectomies	266 (9.2%)
# left hepatectomies	343 (11.9%)
# left lateral sectionectomy	420 (14.5%)
# living related donors	18 (0.6%)
# ALPPS	24 (0.8%)
% conversion	9.7% (0–60)
% morbidity	13% (0–50)
% mortality	0.7% (0–5)

Regarding the indication for liver resection, 1241 cases (43%) were done for benign diseases and 1646 (57%) for malignant diseases. From the malignant group, 62% were for colorectal liver metastasis and 20% for hepatocellular carcinoma. The majority of resections (76.5%) were minor hepatectomies, and 678 resections (23.5%) were major hepatectomies. There were 266 right hepatectomies, 343 left hepatectomies, 420 left lateral sectionectomies, and 24 ALPPS.

The percentage of LLR over open resections was, in 57% of the centers, between 10 and 29% (Fig. 2). In only three groups, more than 70% of resections were done by minimally invasive approach.

Regarding the perioperative results, conversion rate was 9.7%; transfusion rate 13%, overall morbidity 13%, and mortality 0.7%. When considering morbidity and mortality according to the volume of operated patients per center, in high-volume centers (more than 100 LLR) the mean morbidity rate was 23% (4.6–45.6%), mean mortality rate was 0.7% (0–21%), and mean conversion rate was 10.9% (2.7–31.4%). In low-volume centers, the mean morbidity rate was 11% (0–50%), mean mortality rate was 0.8% (0–5%), and mean conversion rate was 9.2% (0–60%).

Discussion

Despite increasingly performed, MILS is employed in a small percentage of liver resections (5–30%), although some expert centers have reported higher rates, reaching 50–80% of liver resections done by laparoscopy [24, 30, 31]. An analysis including liver resections

Fig. 1 Total number of laparoscopic liver resections per group in South America





Fig. 2 Percentage of laparoscopic liver resections over open liver resections by group



performed in France along the year of 2013 showed that only 15% of liver resections were performed through minimally invasive approach [32]. Data from North American centers indicate that less than 10% of all liver resections done for benign conditions were done by laparoscopy [33]. The majority of reported cases are minor resections; however, the proportion of major liver resections is increasing in the last years [30].

In fact, there are few centers worldwide with extensive experience with LLR [30]. A recent survey involving 27 hepatobiliary expert centers worldwide showed different numbers according to the region: the proportion of LLR was 34.6% in Asia-Pacific, 35.6% in Europe, and 27.4% in North/South America [34]. Our objective was to map the use of MILS in South America.

In addition to the aforementioned issues regarding MILS acceptance, in developing countries there is still a barrier due to high equipment costs and difficult access to technology. In a meta-analysis by Limongelli et al., LLR was associated with lower ward stay cost than OLR (2972 USD vs. 5291 USD), but costs related to operation (equipment and theater) were higher in the group of patients undergoing LLR [35]. Despite these difficulties, LLR has spread

widely throughout all South America. In a recent survey in Brazil, more than 90% of liver surgery groups perform LLR [36]. In the present survey, all groups that answered the survey perform LLR, and in the majority of groups (57%), the proportion of LLR over all liver resections ranged between 10 and 29%. Aldrighetti et al., in an Italian national survey on LLR, reported that the proportion of minimally invasive over the total number of liver resections was 10.3% of all resections, with a significant variety between ranging from 0.9 to 58%, with a median of 11% [37].

Indications for resection of malignant tumors in recent years have trespassed the indications for benign diseases, figures observed in the literature [22, 37] and in our study. The reasons for this inversion were based on data demonstrating the oncological effectiveness of MILS both in colorectal cancer metastasis and hepatocellular carcinoma [28].

One of the initial concerns in LLR, especially in major liver resections, was the risk of gas embolism. The incidence is low ranging between 0.2 and 1.45%; and no influence on postoperative morbidity and mortality was

reported [38, 39]. In this series, no cases of embolism were observed.

The major operative concern regarding LLR is massive intraoperative bleeding. Reported blood loss varies between series and is directly related to the type and difficulty of LLR [40–46]. In several meta-analyses, intraoperative bleeding tends to be lower with the laparoscopic approach than OLR resulting in decreased requirement for blood transfusions [28, 47–49]. In a recent review, 8.3% of the patients submitted to LLR received blood transfusion [22], and similar numbers were observed in the present survey where the transfusion rate was 13%.

In a large meta-analyses, Ciria et al. pointed out the safety of LLR showing that in minor-only resection series and in combined series, the overall rate of complications, estimated blood loss, rate of transfusions, and hospital stay are lower in laparoscopic group. In addition, in major-only resection series, the overall rate of complications, estimated blood loss, and hospital stay favor the minimally invasive approach [22].

The reported conversion rate ranges from 0 to 20% [14], varying mostly according to the indication and type of LLR. However, with the acquirement of surgical expertise the conversion rate can be reduced to less than 5% in expert centers [28, 50, 51]. In the present series, the 9.7% of conversion rate can be explained by the high number of groups (52.9%) with less than 50 LLR. Similarly, an Italian multicentric survey with 1677 patients showed a conversion rate of 10.7% [37].

Regarding the complexity of laparoscopic liver resections, in the South American data, 23.5% of the cases were major liver resections, numbers coincident with the world experience (24.2%) [22] showing the gain of experience and expertise of Latin groups. Of interest, is the large number of laparoscopic ALPPS procedures done in South

America, numbers explained by the great enthusiasm with this new technique by groups from Argentina and Brazil [52–54].

In large series, overall morbidity rate ranged from 3.2 [55] to 45%. Jackson et al. [49] analyzing 47 studies showed that patients who underwent LLR had lower postoperative complication rates when compared with open surgery. In a recent prospective trial comparing open and laparoscopic resection for colorectal liver metastasis, the postoperative complication rate was 19% in the laparoscopic group and 31% in the open surgery group [23]. In the present survey, postoperative complication rate was low (13%), probably because a possible selection bias for the minimally invasive procedure, which implies in a large number of minor resections (76.5%). Moreover, low-volume centers presented lower morbidity and conversion rates when compared to high-volume ones; this can also be explained by the higher number of minor and non-complex resections performed in low-volume centers.

Recent series from high-volume centers report mortality for LLR ranging from 0 to 2.4% [6, 20, 42, 44, 46, 50, 51, 56–66]. Jackson et al. pooled the results of 40 studies comparing mortality rates between LLR and OLR and found no significant difference in mortality between groups [49]. In fact, most individual-based studies failed to detect mortality differences between laparoscopic and open resection. However, significant lower 90-day mortality rates (1.3% vs. 5.8%, $P = 0.006$) in patients undergoing laparoscopic left lateral sectionectomy for primary liver malignancy were reported in a population-based study after adjustment for confounding [67]. In a recent review, mortality rate for LLR was 0.4% [22], and in the present series, 0.7%. This data is similar than in the Italian national survey where the mortality rate was 0.2% [36]. These excellent results could be explained by a possible selection bias in the LLR group; moreover, the complex resections were mostly performed in groups of expert surgeons.

A consensual observation is that LLR should be performed by surgeons with extensive training in hepatobiliary and advanced laparoscopic surgery [41]. Therefore, fellowships in specialized centers should offer high-level training in order to accomplish competence in both domains. Indeed, in the Morioka expert consensus in 2014, one of the conclusions of the working group was the need for a formal structure of education for those interested in performing major laparoscopic LLR [24]. In South America, besides some groups with large experience in LLR, there are training centers for minimally invasive liver surgery in Argentina (two), Brazil (three), and Chile (one).

LLR has evolved greatly over the past two decades and, despite initial difficulties and even skepticism, the excellent reported results made the minimally invasive approach

Table 2 Total number of laparoscopic liver resections (LLR) performed per country or region and the relative number LLR/millions of inhabitants (population) in different places of the world

Place	# LLR	Population	# LLRs/million
Japan	20,000*	126,746,000	157.8
France	4509**	66,990,000	67.3
Italy	3302****	60,484,000	54.6
USA	5000****	329,970,000	15.1
South America	2887*****	422,500,000	6.8

*Estimated (Go Wakabayashi) first case 1993

**From 2000 to June 2018 (E. Vibert)

***From 2014 to 2019 (N. Rusonniello)

****Estimated (Joe Buell)

*****From 1997 to April 2019 (J. Pekolj)

to be incorporated to the day-by-day surgical practice. Increased experience, especially in specialized centers, allowed more complex surgeries to be performed. The dissemination and interest in LLR were due to the technical challenge and excellent perioperative results. The visits of surgeons beginning their experience in minimally invasive liver surgery to centers of great expertise, in addition to the proliferation of training centers, enabled a great evolution of LLR worldwide.

This survey showed that South American hepatobiliary centers as well as Asian, European, and North American centers have incorporated LLR into their routine (Table 2). Nowadays, worldwide experience counts more than 35,000 cases of LLR and South American centers have made significant contribution for this scenario. Ours results showed a great number of centers performing LLR, dissemination of training centers, and favorable results with low perioperative morbidity and mortality, in accordance with other expert centers around the world.

References

- Cherqui D, Husson E, Hammoud R et al (2000) Laparoscopic liver resections: a feasibility study in 30 patients. *Ann Surg* 232:753–762
- Viganò L, Tayar C, Laurent A et al (2009) Laparoscopic liver resection: a systematic review. *J Hepatobiliary Pancreat Surg* 16:410–421
- Cherqui D (2003) Laparoscopic liver resection. *Br J Surg* 90:644–646
- D'Albuquerque LA, Herman P (2006) Laparoscopic hepatectomy: is it a reality? *Arq Gastroenterol* 43:243–246
- Abu Hilal M, Di Fabio F, Abu Salameh M et al (2012) Oncological efficiency analysis of laparoscopic liver resection for primary and metastatic cancer: a single-center UK experience. *Arch Surg* 147:42–48
- Herman P, Coelho FF, Perini MV et al (2012) Hepatocellular adenoma: an excellent indication for laparoscopic liver resection. *HPB (Oxford)* 14:390–395
- Doughtie CA, Egger ME, Cannon RM et al (2013) Laparoscopic hepatectomy is a safe and effective approach for resecting large colorectal liver metastases. *Am Surg* 79:566–571
- Cauchy F, Schwarz L, Scatton O et al (2014) Laparoscopic liver resection for living donation: where do we stand? *World J Gastroenterol* 20:15590–15598
- Scatton O, Katsanos G, Boillot O et al (2015) Pure laparoscopic left lateral sectionectomy in living donors: from innovation to development in France. *Ann Surg* 261:506–512
- Luo LX, Yu ZY, Bai YN (2014) Laparoscopic hepatectomy for liver metastases from colorectal cancer: a meta-analysis. *J Laparoendosc Adv Surg Tech A* 24:213–222
- Schiffman SC, Kim KH, Tsung A et al (2015) Laparoscopic versus open liver resection for metastatic colorectal cancer: a meta-analysis of 610 patients. *Surgery* 157:211–222
- Xiong JJ, Altaf K, Javed MA et al (2012) Meta-analysis of laparoscopic vs open liver resection for hepatocellular carcinoma. *World J Gastroenterol* 18:6657–6668
- Twajj A, Pucher PH, Sodergren MH et al (2014) Laparoscopic vs open approach to resection of hepatocellular carcinoma in patients with known cirrhosis: systematic review and meta-analysis. *World J Gastroenterol* 20:8274–8281
- Li N, Wu YR, Wu B et al (2012) Surgical and oncologic outcomes following laparoscopic versus open liver resection for hepatocellular carcinoma: a meta-analysis. *Hepatol Res* 42:51–59
- Yin Z, Fan X, Ye H et al (2013) Short- and long-term outcomes after laparoscopic and open hepatectomy for hepatocellular carcinoma: a global systematic review and meta-analysis. *Ann Surg Oncol* 20:1203–1215
- Ye SP, Qiu H, Liao SJ et al (2019) Mini-invasive vs open resection of colorectal cancer and liver metastases: a meta-analysis. *World J Gastroenterol* 25(22):2819–2832
- Ciria R, Ocaña S, Gomez-Luque I et al (2020) A systematic review and meta-analysis comparing the short- and long-term outcomes for laparoscopic and open liver resections for liver metastases from colorectal cancer. *Surg Endosc* 34(1):349–360
- Jayaraman S, Khakhar A, Yang H, Bainbridge D, Quan D (2009) The association between central venous pressure, pneumoperitoneum, and venous carbon dioxide embolism in laparoscopic hepatectomy. *Surg Endosc* 23:2369–2373. <https://doi.org/10.1007/s00464-009-0359-9>
- Gaillard M, Tranchart H, Dagher I (2014) Laparoscopic liver resections for hepatocellular carcinoma: current role and limitations. *World J Gastroenterol* 20:4892–4899
- Ding G, Cai W, Qin M (2015) Pure laparoscopic versus open liver resection in treatment of hepatolithiasis within the left lobes: a randomized trial study. *Surg Laparosc Endosc Percutan Tech* 25:392–394
- Tranchart H, Dagher I (2014) Laparoscopic liver resection: a review. *J Visc Surg* 151:107–115
- Ciria R, Cherqui D, Geller DA et al (2016) Comparative short-term benefits of laparoscopic liver resection: 9000 cases and climbing. *Ann Surg* 263(4):761–777
- Fretland AA, Dagenborg VJ, Bjørnelv GMW et al (2018) Laparoscopic versus open resection for colorectal liver metastases: the OSLO-COMET randomized controlled trial. *Ann Surg* 267(2):199–207
- Wakabayashi G, Cherqui D, Geller DA et al (2015) Recommendations for laparoscopic liver resection: a report from the second international consensus conference held in Morioka. *Ann Surg* 261:619–629
- Kalil AN, Mastalir ET (2002) Laparoscopic hepatectomy for benign liver tumors. *Hepatogastroenterology* 49(45):803–805
- Claria RS, Ardiles V, Palavecino ME et al (2009) Laparoscopic resection for liver tumors: initial experience in a single center. *Surg Laparosc Endosc Percutan Tech* 19(5):388–391
- Machado MA, Surjan RC, Basseres T et al (2016) The laparoscopic Glissonian approach is safe and efficient when compared with standard laparoscopic liver resection: results of an observational study over 7 years. *Surgery* 160(3):643–651
- Coelho FF, Kruger JA, Fonseca GM et al (2016) Laparoscopic liver Resection: experience Based Guidelines. *World J Gastrointest Surg* 8(1):5–26
- Herman P, Kruger JAP, Perini MV et al (2013) Laparoscopic hepatic posterior sectionectomy: a hand assisted approach. *Ann Surg Oncol* 20(4):1266
- Dagher I, Gayet B, Tzanis D et al (2014) International experience for laparoscopic major liver resection. *J Hepatobiliary Pancreat Sci* 21:732–736
- Cannon RM, Saggi B, Buell JF (2014) Evaluation of a laparoscopic liver resection in the setting of cirrhosis. *HPB (Oxford)* 16:164–169

32. Farges O, Goutte N, Dokmak S et al (2014) How surgical technology translates into practice: the model of laparoscopic liver resections performed in France. *Ann Surg* 260:916–921
33. Kim Y, Amini N, He J et al (2015) National trends in the use of surgery for benign hepatic tumors in the United States. *Surgery* 157:1055–1064
34. Kawaguchi Y, Hasegawa K, Wakabayashi G et al (2016) Survey results on daily practice in open and laparoscopic liver resections from 27 centers participating in the second International Consensus Conference. *JHPB Sci* 23(5):283–288
35. Limongelli P, Vitiello C, Belli A et al (2014) Costs of laparoscopic and open liver and pancreatic resection: a systematic review. *World J Gastroenterol* 20:17595–17602
36. Fonseca GM, Jeismann VB, Kruger JAP et al (2018) Liver resection in BRAZIL: a national survey. *Arq Bras Cir Dig* 31(1):e1355
37. Aldrighetti L, Belli G, Boni L et al (2015) Italian experience in minimally invasive liver surgery: a national survey. *Updates Surg* 67:129–140
38. Dagher I, O'Rourke N, Geller DA et al (2009) Laparoscopic major hepatectomy: an evolution in standard of care. *Ann Surg* 250:856–860
39. Otsuka Y, Katagiri T, Ishii J et al (2013) Gas embolism in laparoscopic hepatectomy: what is the optimal pneumoperitoneal pressure for laparoscopic major hepatectomy? *J Hepatobiliary Pancreat Sci* 20:137–140
40. Koffron AJ, Auffenberg G, Kung R et al (2007) Evaluation of 300 minimally invasive liver resections at a single institution: less is more. *Ann Surg* 246:385–392
41. Buell JF, Cherqui D, Geller DA et al (2009) The international position on laparoscopic liver surgery: the Louisville Statement 2008. *Ann Surg* 250:825–830
42. Cai X, Li Z, Zhang Y et al (2014) Laparoscopic liver resection and the learning curve: a 14-year, single-center experience. *Surg Endosc* 28:1334–1341
43. Gobardhan PD, Subar D, Gayet B (2014) Laparoscopic liver surgery: an overview of the literature and experiences of a single centre. *Best Pract Res Clin Gastroenterol* 28:111–121
44. Troisi RI, Montalti R, Van Limmen JG et al (2014) Risk factors and management of conversions to an open approach in laparoscopic liver resection: analysis of 265 consecutive cases. *HPB (Oxford)* 16:75–82
45. Bryant R, Laurent A, Tayar C et al (2009) Laparoscopic liver resection—understanding its role in current practice: the Henri Mondor Hospital experience. *Ann Surg* 250:103–111
46. Buell JF, Thomas MT, Rudich S et al (2008) Experience with more than 500 minimally invasive hepatic procedures. *Ann Surg* 248:475–486
47. Mirnezami R, Mirnezami AH, Chandrakumaran K et al (2011) Short- and long-term outcomes after laparoscopic and open hepatic resection: systematic review and meta-analysis. *HPB (Oxford)* 13:295–308
48. Simillis C, Constantinides VA, Tekkis PP et al (2007) Laparoscopic versus open hepatic resections for benign and malignant neoplasms: a meta-analysis. *Surgery* 141:203–211
49. Jackson NR, Hauch A, Hu T et al (2015) The safety and efficacy of approaches to liver resection: a meta-analysis. *JSL* 19(e2014):00186
50. Abu Hilal M, Di Fabio F, Teng MJ et al (2011) Surgical management of benign and indeterminate hepatic lesions in the era of laparoscopic liver surgery. *Dig Surg* 28:232–236
51. Hasegawa Y, Nitta H, Sasaki A et al (2015) Long-term outcomes of laparoscopic versus open liver resection for liver metastases from colorectal cancer: a comparative analysis of 168 consecutive cases at a single center. *Surgery* 157:1065–1072
52. Pekolj J, Alvarez FA, Biagiola D et al (2018) Totally laparoscopic mini-ALPPS using a novel approach of laparoscopic-assisted transmesenteric portal vein embolization. *J Laparoendoscopic Adv Surg Tech* 28(10):1229–1233
53. Machado MA, Makdissi FF, Surjan RC et al (2017) Transition from open to laparoscopic ALPPS for patients with very small FLR: the initial experience. *HPB (Oxford)* 19(1):59–66
54. Schelotto PB, Gondolesi G (2017) Laparoscopy in ALPPS procedure: when we can do it? *Ann Surg* 265(4):e30–e31
55. Cai XJ, Yu H, Liang X et al (2006) Laparoscopic hepatectomy by curettage and aspiration. Experiences of 62 cases. *Surg Endosc* 20:1531–1535
56. Dokmak S, Raut V, Aussilhou B et al (2014) Laparoscopic left lateral resection is the gold standard for benign liver lesions: a case–control study. *HPB (Oxford)* 16:183–187
57. Lee KF, Chong CN, Wong J et al (2011) Long term results of laparoscopic hepatectomy versus open hepatectomy for hepatocellular carcinoma: a case-matched analysis. *World J Surg* 35:2268–2274
58. Cheung TT, Poon RT, Yuen WK et al (2013) Long-term survival analysis of pure laparoscopic versus open hepatectomy for hepatocellular carcinoma in patients with cirrhosis: a single-center experience. *Ann Surg* 257:506–511
59. Memeo R, de Angelis N, Compagnon P et al (2014) Laparoscopic vs. open liver resection for hepatocellular carcinoma of cirrhotic liver: a case–control study. *World J Surg* 38:2919–2926
60. Yamashita Y, Ikeda T, Kurihara T et al (2014) Long-term favorable surgical results of laparoscopic hepatic resection for hepatocellular carcinoma in patients with cirrhosis: a single-center experience over a 10-year period. *J Am Coll Surg* 219:1117–1123
61. Martin RC, Mbah NA, St Hill R et al (2015) Laparoscopic versus open hepatic resection for hepatocellular carcinoma: improvement in outcomes and similar cost. *World J Surg* 39:1519–1526
62. Lee JJ, Conneely JB, Smoot RL et al (2015) Laparoscopic versus open liver resection for hepatocellular carcinoma at a North-American Centre: a 2-to-1 matched pair analysis. *HPB (Oxford)* 17:304–310
63. Han HS, Shehta A, Ahn S et al (2015) Laparoscopic versus open liver resection for hepatocellular carcinoma: case matched study with propensity score matching. *J Hepatol* 15(63):643–650
64. Cannon RM, Scoggins CR, Callender GG et al (2012) Laparoscopic versus open resection of hepatic colorectal metastases. *Surgery* 152:567–573
65. Kubota Y, Otsuka Y, Tsuchiya M et al (2014) Efficacy of laparoscopic liver resection in colorectal liver metastases and the influence of preoperative chemotherapy. *World J Surg Oncol* 12:351
66. de'Angelis N, Eshkenazy R, Brunetti F et al (2015) Laparoscopic versus open resection for colorectal liver metastases: a single-center study with propensity score analysis. *J Laparoendoscopic Adv Surg Tech A* 25:12–20
67. Goutte N, Bendersky N, Barbier L et al (2017) Laparoscopic left lateral sectionectomy: a population-based study. *HPB* 19(2):118–125

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.